

Genera of zero-divisor graphs with respect to ideals

Hsin-Ju Wang *

Department of Mathematics, National Chung Cheng University,
Chiayi 621, Taiwan

Abstract. In this paper, we study the genera of zero-divisor graphs with respect to ideals in finite rings.

1 Introduction

In this paper, R will denote a commutative ring with 1. Often, but not always, we will also assume that R is finite. If S is a subset of R , we denote $S - \{0\}$ by S^* . Also, we use $\mathbb{N}$ for the natural numbers and $\mathbb{F}_q$ for the finite field of q elements.

A subject of study linking commutative ring theory with graph theory has been the concept of the *zero-divisor graph* of a commutative ring. Let R be a ring. The zero-divisor graph of R , denoted $\Gamma(R)$, is a simple graph whose vertices are the nonzero zero-divisors of R with two distinct vertices x and y joined by an edge if and only if $xy = 0$. This definition was introduced by Anderson and Livingston in [4] and later was studied extensively in [1], [2], [3], [4], [9] and [10]. Recently, Redmond extends the concept of zero-divisor graphs to *zero-divisor graphs with respect to ideals*. Namely, a simple graph $\Gamma_I(R)$ with vertices $\{x \in R - I \mid xy \in I \text{ for some } y \in R - I\}$ in which distinct vertices x and y are adjacent if and only if $xy \in I$.

One of the primitive subjects of topological graph theory is to embed a graph into a surface. In plain words, it is to draw a graph on a surface so that there is no crossing for any two edges. One simple question which one may ask is that "What kind of rings can have $\gamma(\Gamma(R)) = g$ (see definition in

*e-mail: hjwang@math.ccu.edu.tw

section 2)?" . For example, in [5], Chiang-Hsieh and Wang find all finite rings that have genus one. When concerning to zero-divisor graphs with respect to ideals, similar question also has been discussed. In [7], Redmond find a sufficient and necessary condition for which $\Gamma_I(R)$ is planar. The goal of this note is to find all finite rings and ideals for which the genus of $\Gamma_I(R)$ is one.

In order to obtain our main results, we review in section 2 some background from graph theory and derive a criterion for a graph to have bigger genus than its subgraph.

In section 3, we establish the relationship between the graph $\Gamma(R/I)^{(t)}$ and the graph $\Gamma_I(R)$ if $|I| = t$, where $G^{(t)}$ is defined in section 2 for any finite simple graph.

In section 4, we first show that if $\omega(\Gamma(R/I)) \geq 4$ or $\gamma(\Gamma(R/I)) \geq 1$, then $\gamma(\Gamma_I(R)) \geq 2$. Then we state and prove some equivalent conditions for which $\gamma(\Gamma_I(R)) = 1$.

2 Preliminary

We review some background from graph theory in this section.

A simple graph G is an ordered pair of disjoint sets (V, E) such that $V = V(G)$ is the vertex set of G and $E = E(G)$ is the edge set of G . The *order* of a graph G , written by $|G|$, is the cardinality of $V(G)$. A *subgraph* of G is a graph having all of its vertices and edges in G . Let $V' \subseteq V(G)$; then $G - V'$ is the subgraph of G obtained by deleting the vertices in V' and all edges incident with them. Similarly, if $E' \subseteq E(G)$, then $G - E'$ is the subgraph of G obtained by deleting the edges in E' . For any set S of vertices of G , we use the symbol $\langle S \rangle$ for the subgraph induced by S .

A *bipartite* graph (bigraph) G is a graph whose point set V can be partitioned into two subsets V_1 and V_2 such that every line of G joins V_1 with V_2 . If G contains every line joining V_1 and V_2 , then G is a *complete bipartite* graph. If $|V_1| = m$ and $|V_2| = n$, we use the symbol $K_{m,n}$ for the complete bipartite graph. Moreover, if $m = 1$ or $n = 1$, then G is called a *star graph*. A graph in which each pair of distinct vertices is joined by an edge is called a *complete graph*. We use K_n for the complete graph with n vertices. For a graph G , the *clique number*, $\omega(G)$, is the greatest integer $n \geq 1$ such that K_n is a subgraph of G . If K_n is a subgraph of G for every n then $\omega(G) = \infty$.

Remark 2.1 In [7, Theorem 7.2], the author states that $\omega(G) \leq 2$ if and

only if G contains no cycles. However, this statement is wrong as if $G = C_4$ (the cycle of length 4) satisfies $\omega(G) \leq 2$.

A simple graph is said to be *planar* if it can be drawn in the plane or on the surface of a sphere. It is known that $K_{3,3}$ and K_5 are not planar and can be drawn without crossings on the surface of a torus. The torus can be thought of as a sphere with one *handle*. More generally, a surface is said to be of genus g if it is topologically homeomorphic to a sphere with g handles. Thus the genus of a sphere is 0 and the one of torus is one. A graph can be drawn without crossings on the surface of genus g , but not on one of genus $g-1$, is called a graph of genus g . We write $\gamma(G)$ for the genus of a graph G . Therefore $\gamma(K_{3,3}) = \gamma(K_5) = 1$. A well-known fact is that if G is a connected graph of genus g , then any presentation of G on a surface of genus g satisfies $n - m + f = 2 - 2g$ and $2m \geq 3f$, where n, m, f are the vertices, edges and faces according to the presentation. There are some known results from [8] which will be used later.

Lemma 2.2 $\gamma(K_n) = \{\frac{1}{12}(n-3)(n-4)\}$, where $\{x\}$ is the least integer that is greater than or equal to x . In particular, $\gamma(K_n) = 1$ if $n = 5, 6, 7$.

Proof. See, for example [8]. $\square$

Lemma 2.3 $\gamma(K_{m,n}) = \{\frac{1}{4}(m-2)(n-2)\}$, where $\{x\}$ is the least integer that is greater than or equal to x . In particular, $\gamma(K_{4,4}) = \gamma(K_{3,n}) = 1$ if $n = 3, 4, 5, 6$.

Proof. See, for example [8]. $\square$

For later use, we introduce the following notion:

Definition 2.4 Let G be a finite simple graph with vertex set $\{w_1, \dots, w_m\}$ and edge set $E(G)$; then $G^{(t)}$ is the finite simple graph with vertex set $\{w_{ij} \mid 1 \leq i \leq t, 1 \leq j \leq m\}$ and edge set $\{w_{ij}w_{kl} \mid j \neq l, w_jw_l \in E(G)\}$.

Remark 2.5 Observe that $w_{ij}w_{kl}$ is not an edge of $G^{(t)}$ if $j = l$.

Example 2.6 If $G = K_{m,n}$ then $G^{(t)} = K_{mt,nt}$. If $G = K_n$ then $G^{(t)} = K_{t,\dots,t}$.

There are some results concerning the genera of $G^{(t)}$.

Proposition 2.7 *Let G be a finite simple graph. Then the following hold:*

- (a) *If K_2 is a subgraph of G , then $\gamma(G^{(5)}) \geq 2$.*
- (b) *If K_5 is a subgraph of G , then $\gamma(G^{(2)}) \geq 2$.*
- (c) *If $K_{1,3}$ is a subgraph of G , then $\gamma(G^{(3)}) \geq 2$.*
- (d) *If $K_{2,3}$ is a subgraph of G , then $\gamma(G^{(2)}) \geq 2$.*

Proof. (a) We may assume that $G = K_2$. Then by Example 2.6, $G^{(5)} = K_{5,5}$. Therefore, $\gamma(G^{(5)}) \geq 2$ by Lemma 2.3.

(b) We may assume that $G = K_5$. Let $V(G) = \{w_1, \dots, w_5\}$ and $E(G) = \{w_i w_j \mid 1 \leq i < j \leq 5\}$. Then $V(G^{(2)}) = \{w_{ij} \mid 1 \leq i \leq 2, 1 \leq j \leq 5\}$ and $E(G^{(2)}) = \{w_{ij} w_{kl} \mid j \neq l\}$. Let $u_1 = w_{11}$, $u_2 = w_{12}$, $u_3 = w_{21}$, $u_4 = w_{22}$, $v_1 = w_{13}$, $v_2 = w_{14}$, $v_3 = w_{15}$, $v_4 = w_{23}$, $v_5 = w_{24}$ and $v_6 = w_{25}$; then $u_i v_j$ is an edge of $G^{(2)}$, so that $K_{4,6}$ is a subgraph of $G^{(2)}$, it follows that $\gamma(G^{(2)}) \geq 2$ by Lemma 2.3.

(c) We may assume that $G = K_{1,3}$. Then by Example 2.6, $G^{(3)} = K_{3,9}$. Therefore, $\gamma(G^{(3)}) \geq 2$ by Lemma 2.3.

(d) We may assume that $G = K_{2,3}$. Then by Example 2.6, $G^{(2)} = K_{4,6}$. Therefore, $\gamma(G^{(2)}) \geq 2$ by Lemma 2.3.

□

To end this section, we illustrate the following useful Lemma.

Lemma 2.8 *Let G be a connected graph and G_1, G_2 be subgraphs of G . Suppose the following hold:*

- (a) *$V(G) = V(G_1) \cup V(G_2)$ and $V(G_1) \cap V(G_2) = \emptyset$.*
 - (b) *$V(G_1) = \{v_1, \dots, v_4\}$ and $G_1 = K_4$.*
 - (c) *For every i , there is an edge in G joins v_i with G_2 .*
- Then $\gamma(G) > \gamma(G_2)$.*

Proof. Let $g = \gamma(G_2)$. Suppose on the contrary that $\gamma(G) = \gamma(G_2)$. Assume that $\{F_1, \dots, F_m\}$ and $\{F'_1, \dots, F'_{m'}\}$ are faces of G and G_2 when drawing G and G_2 on a surface S_g of genus g . Then $\{F'_1, \dots, F'_{m'}\}$ can be obtained by deleting $v_1, \dots, v_4$ and all edges incident with $v_1, \dots, v_4$ from the presentation of G . Further, $\{F_1, \dots, F_m\}$ can be recovered by inserting $v_1, \dots, v_4$ and all edges incident with $v_1, \dots, v_4$ into the presentation corresponding to G_2 . Let F'_{t_i} denote the face of G_2 into which v_i is inserted during the recovering process from G_2 to G . We note that $v_i v_j \in E(G)$ for every $i \neq j$; therefore

all v_i should be inserted into the same face, say $F = F'_1$, of G_2 to avoid any crossings, i.e., $t_1 = \dots = t_4 = 1$. Let e_i be an edge join v_i to G_2 for $i = 1, \dots, 4$. After inserting G_1 into F we obtain Figure 1. However, it is easy to see from Figure 1 that we can not insert $e_1, \dots, e_4$ into F without crossings, a contradiction. Therefore, we conclude that $\gamma(G) > \gamma(G_2)$. $\square$




Figure 1

3 Basis properties of $\Gamma_I(R)$

Throughout, let R be a commutative ring with 1 and I be an ideal of R . We use $\Gamma(R)$ to denote the zero-divisor graph of R . In [7], Redmond extends the concept of zero-divisor graphs as follows:

Definition 3.1 *Let R be a commutative ring and let I be an ideal of R . We define a simple graph $\Gamma_I(R)$ with vertices $\{x \in R - I \mid xy \in I \text{ for some } y \in R - I\}$, where distinct vertices x and y are adjacent if and only if $xy \in I$.*




Figure 2-1




Figure 2-2

Example 3.2 In Figure 2-1, $R = \mathbb{Z}_6 \times \mathbb{Z}_2$ and $I = 0 \times \mathbb{Z}_2$. In Figure 2-2, $R = \mathbb{Z}_8 \times \mathbb{Z}_2$ and $I = 0 \times \mathbb{Z}_2$.

Remark 3.3 (a) By Definition 3.1, if $\{x_\lambda + I \mid \lambda \in \Lambda\}$ is the set of nonzero zero-divisors of R/I , then the vertex set of $\Gamma_I(R)$ is $\{x_\lambda + a \mid \lambda \in \Lambda, a \in I\}$, therefore $|\Gamma_I(R)| = |I| \cdot |\Gamma(R/I)|$.

(b) By Definition 2.4 and Definition 3.1, if $|I| = t$ then $\Gamma(R/I)^{(t)}$ is a subgraph of $\Gamma_I(R)$. More precisely, if $\{x_1 + I, \dots, x_m + I\}$ is the set of nonzero zero-divisors of R/I , $I = \{a_1, \dots, a_t\}$ and $G = \Gamma(R/I)$, then the vertex set of $\Gamma_I(R)$ is $\{x_j + a_i \mid 1 \leq i \leq t, 1 \leq j \leq m\}$ and the edge set of $\Gamma_I(R)$ is $\{(x_j + a_i)(x_l + a_k) \mid x_j x_l \in I\}$. However, the edge set of $G^{(t)}$ is $\{(x_j + a_i)(x_l + a_k) \mid j \neq l, x_j x_l \in I\}$. Thus, $\Gamma(R/I)^{(t)}$ is a subgraph of $\Gamma_I(R)$. Also, from this one can easily see that $\Gamma(R/I)^{(t)} = \Gamma_I(R)$ if and only if I is radical.

(c) From the above one can see that $\Gamma_I(R)$ is depend only on R/I and $|I|$. Therefore the graph $\Gamma_I(R)$ and the graph $\Gamma_{0 \times A}(R/I \times A)$ are the same, where A is any ring with $|A| = |I|$.

We list some properties of $\Gamma_I(R)$ in the following. Part of them come from [6] and [7].

Theorem 3.4 $\Gamma_I(R)$ is finite if and only if R is finite. Moreover, $\Gamma_I(R)$ is connected.

Theorem 3.5 Let R be a finite ring and I be a proper nonzero ideal of R . Then the diameter of $\Gamma_I(R)$ is less than or equal to 3. If $\Gamma_I(R)$ contains a cycle then $gr(\Gamma_I(R)) \leq 4$.

Theorem 3.6 Let I be a finite ideal of a ring R and let J be a finite ideal of a ring S such that $\sqrt{I} = I$ and $\sqrt{J} = J$. Then the following hold:

(a) If $|I| = |J|$ and $\Gamma(R/I) \cong \Gamma(S/J)$, then $\Gamma_I(R) \cong \Gamma_J(S)$.

(b) If $\Gamma_I(R) \cong \Gamma_J(S)$, then $\Gamma(R/I) \cong \Gamma(S/J)$

Theorem 3.7 Let I be a radical ideal of a ring R such that $I = \cap_{1 \leq i \leq t} \mathfrak{p}_i$ is a minimal primary decomposition, where $\mathfrak{p}$ is a prime ideal for every i . Then $\omega(\Gamma_I(R)) = n$.

4 The genera of $\Gamma_I(R)$

In this section, we find some sufficient and necessary conditions for a finite ring R and a nonzero ideal I of R to have $\gamma(\Gamma_I(R)) \leq 1$. To begin with, we state some known results and list some easy observations:

Theorem 4.1 [7, Theorem 7.2] *Let R be a finite ring and I be a proper nonzero ideal of R which is not a prime ideal. Then $\Gamma_I(R)$ is planar if and only if $\Gamma(R/I)$ contains no cycles and either (a) $|I| = 2$ or (b) $|\Gamma(R/I)| = 1$ and $|I| \leq 4$.*

Theorem 4.2 [5] *Let $(R, \mathfrak{m})$ be a finite local ring which is not a field. Then $\Gamma(R)$ is planar if and only if R is isomorphic to one of the following 29 rings.*

$$\begin{aligned} & \mathbb{Z}_4, \mathbb{Z}_8, \mathbb{Z}_9, \mathbb{Z}_{16}, \mathbb{Z}_{25}, \mathbb{Z}_{27}, \frac{\mathbb{Z}_2[x]}{(x^2)}, \frac{\mathbb{Z}_2[x]}{(x^3)}, \frac{\mathbb{Z}_2[x]}{(x^4)}, \frac{\mathbb{Z}_2[x, y]}{(x^2, xy, y^2)}, \frac{\mathbb{Z}_2[x, y]}{(x^2, y^2)}, \frac{\mathbb{Z}_2[x, y]}{(x^3, xy, y^2 - x^2)}, \\ & \frac{\mathbb{F}_4[x]}{(x^2)}, \frac{\mathbb{Z}_3[x]}{(x^2)}, \frac{\mathbb{Z}_3[x]}{(x^3)}, \frac{\mathbb{Z}_4[x]}{(x^2)}, \frac{\mathbb{Z}_4[x]}{(x^2 + x + 1)}, \frac{\mathbb{Z}_4[x]}{(2x, x^2)}, \frac{\mathbb{Z}_4[x]}{(x^2 - 2, x^4)}, \frac{\mathbb{Z}_4[x]}{(x^3 - 2, x^4)}, \\ & \frac{\mathbb{Z}_4[x]}{(x^2 - 2, x^3)}, \frac{\mathbb{Z}_4[x]}{(x^3, x^2 - 2x)}, \frac{\mathbb{Z}_4[x]}{(x^3 + x^2 - 2, x^4)}, \frac{\mathbb{Z}_4[x, y]}{(x^2, y^2, xy - 2)}, \frac{\mathbb{Z}_4[x, y]}{(x^3, x^2 - 2, xy, y^2 - 2)}, \\ & \frac{\mathbb{Z}_5[x]}{(x^2)}, \frac{\mathbb{Z}_8[x]}{(x^2 - 4, 2x)}, \frac{\mathbb{Z}_9[x]}{(x^2 - 3, x^3)}, \frac{\mathbb{Z}_9[x]}{(x^2 + 3, x^3)}. \end{aligned}$$

Theorem 4.3 [5] *Let $(R, \mathfrak{m})$ be a finite local ring which is not a field. Then $\gamma(\Gamma(R)) = 1$ if and only if R is isomorphic to one of the following 18 rings.*

$$\begin{aligned} & \mathbb{Z}_{32}, \mathbb{Z}_{49}, \frac{\mathbb{Z}_2[x]}{(x^5)}, \frac{\mathbb{F}_8[x]}{(x^2)}, \frac{\mathbb{Z}_2[x, y]}{(x^3, xy, y^2)}, \frac{\mathbb{Z}_2[x, y, z]}{(x, y, z)^2}, \frac{\mathbb{Z}_4[x]}{(x^3 + x + 1)}, \frac{\mathbb{Z}_4[x]}{(x^3 - x + 1)}, \\ & \frac{\mathbb{Z}_4[x]}{(x^3 - 2, x^5)}, \frac{\mathbb{Z}_4[x]}{(x^4 - 2, x^5)}, \frac{\mathbb{Z}_4[x]}{(x^4 + x^3 - 2, x^5)}, \frac{\mathbb{Z}_4[x]}{(x^3, 2x)}, \frac{\mathbb{Z}_4[x, y]}{(x^3, x^2 - 2, xy, y^2)}, \frac{\mathbb{Z}_4[x, y]}{(2x, 2y, x^2, xy, y^2)}, \\ & \frac{\mathbb{Z}_7[x]}{(x^2)}, \frac{\mathbb{Z}_8[x]}{(x^2, 2x)}, \frac{\mathbb{Z}_8[x]}{(x^2 - 2, x^5)}, \frac{\mathbb{Z}_8[x]}{(3x^2 - 2, x^5)}. \end{aligned}$$

Theorem 4.4 [5] *Let R be a finite ring which has exactly two maximal ideals; then $\Gamma(R)$ is planar if and only if R is isomorphic to one of the following 15 rings.*

$$\begin{aligned} & \mathbb{Z}_2 \times \mathbb{F}_q, \mathbb{Z}_3 \times \mathbb{F}_q, \mathbb{Z}_2 \times \mathbb{Z}_9, \mathbb{Z}_2 \times \frac{\mathbb{Z}_3[x]}{(x^2)}, \mathbb{Z}_2 \times \mathbb{Z}_4, \mathbb{Z}_2 \times \frac{\mathbb{Z}_2[x]}{(x^2)}, \mathbb{Z}_2 \times \frac{\mathbb{Z}_2[x]}{(x^3)}, \\ & \mathbb{Z}_2 \times \frac{\mathbb{Z}_4[x]}{(x^2 - 2, x^3)}, \mathbb{Z}_2 \times \mathbb{Z}_8, \mathbb{Z}_3 \times \mathbb{Z}_9, \mathbb{Z}_3 \times \frac{\mathbb{Z}_3[x]}{(x^2)}, \mathbb{Z}_3 \times \mathbb{Z}_4, \mathbb{Z}_3 \times \frac{\mathbb{Z}_2[x]}{(x^2)}, \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2, \\ & \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_3. \end{aligned}$$

Lemma 4.5 *If $(R, \mathfrak{m})$ is finite local and k is the smallest integer for which $\mathfrak{m}^k = 0$, then $|\mathfrak{m}^i| = t^{n_i} |\mathfrak{m}^{i+1}|$ for $i = 0, \dots, k-1$ if $t = |R/\mathfrak{m}|$. In particular, $|R| = t^n$ for some n .*

Proof. The assertion follows from the fact that $\mathfrak{m}^i/\mathfrak{m}^{i+1}$ is a nonzero vector space over $R/\mathfrak{m}$ for $i = 0, \dots, k-1$ and $|R/\mathfrak{m}| = t$. $\square$

Lemma 4.6 *Let $(R, \mathfrak{m})$ be a finite local ring with $\mathfrak{m}^2 \neq 0$ and $|R/\mathfrak{m}| \geq 3$. Then $\gamma(\Gamma(R)^{(2)}) \geq 2$.*

Proof. Let k be the smallest integer for which $\mathfrak{m}^k = 0$; then $k \geq 3$ by assumption. Let $t = |R/\mathfrak{m}|$; then $t \geq 3$ by assumption. Observe that $|\mathfrak{m}^{k-1} - \{0\}| \geq t-1 \geq 2$ and $|\mathfrak{m}^{k-2} - \mathfrak{m}^{k-1}| \geq (t-1)|\mathfrak{m}^{k-2}| \geq 6$ by Lemma 4.5. Therefore there are distinct elements $u_1, u_2 \in \mathfrak{m}^{k-1} - \{0\}$ and distinct elements $v_1, \dots, v_6 \in \mathfrak{m}^{k-2} - \mathfrak{m}^{k-1}$. Since $k \geq 3$, $u_i v_j = 0$ for every i, j . Thus $K_{2,3}$ is a subgraph of $\Gamma(R)$ and then $\gamma(\Gamma(R)^{(2)}) \geq 2$ by Proposition 2.7(d). $\square$

Lemma 4.7 *Let $(R, \mathfrak{m})$ be a finite local ring such that $\mathfrak{m}^2 \neq 0$. If $\Gamma(R)$ contains no cycles, then $|R/\mathfrak{m}| = 2$.*

Proof. If $|R/\mathfrak{m}| \geq 3$, then from the proof of Lemma 4.6 $u_1 - u_2 - v_1 - u_1$ is a triangle, a contradiction. Thus, $|R/\mathfrak{m}| = 2$. $\square$

Lemma 4.8 *Let $R \cong \mathbb{Z}_2 \times S$, where S is a finite local ring. Then the following hold:*

(a) *If $|\Gamma(S)| \leq 1$, then $\Gamma(R)$ is planar and contains no cycles.*

(b) *If $|\Gamma(S)| \geq 2$, then K_3 and $K_{2,3}$ are subgraphs of $\Gamma(R)$.*

Proof. (a) If S is a finite field, then $\Gamma(R)$ is of course planar and contains no cycles. By Theorem 4.2, if $|\Gamma(S)| = 1$, then $S \cong \mathbb{Z}_4$ or $S \cong \frac{\mathbb{Z}_2[x]}{(x^2)}$, so that $R \cong \mathbb{Z}_2 \times \mathbb{Z}_4$ or $R \cong \mathbb{Z}_2 \times \frac{\mathbb{Z}_2[x]}{(x^2)}$, it follows that $\Gamma(R)$ are trees by [7, Figure 11].

(b) Suppose that $|\Gamma(S)| = 2$. Then $S \cong \mathbb{Z}_9$ or $S \cong \frac{\mathbb{Z}_3[x]}{(x^2)}$ by Theorem 4.2. In either cases, there are two distinct nonzero zero-divisors $a, b \in S$ such that $ab = a^2 = b^2 = 0$. Let $u_1 = (0, a)$, $u_2 = (0, b)$, $v_1 = (1, 0)$, $v_2 = (1, a)$ and $v_3 = (1, b)$; then $\{u_1, u_2, v_1, v_2, v_3\}$ are part of a vertex set of $\Gamma(R)$ such that $u_i v_j = 0$ for all i, j , so that $K_{2,3}$ is a subgraph of $\Gamma(R)$. Moreover, $u_1 - u_2 - v_1 - u_1$ is a triangle contained in $\Gamma(R)$.

Finally assume that $|\Gamma(S)| \geq 3$. Then there are three distinct nonzero zero-divisors $a, b, c \in S$ such that $ab = ac = 0$. Let $u_1 = (0, b)$, $u_2 = (0, c)$, $v_1 = (1, 0)$, $v_2 = (0, a)$ and $v_3 = (1, a)$; then $\{u_1, u_2, v_1, v_2, v_3\}$ are part of a vertex set of $\Gamma(R)$ such that $u_i v_j = 0$ for all i, j , so that $K_{2,3}$ is a subgraph of $\Gamma(R)$. Moreover, $v_1 - v_2 - u_1 - v_1$ is a triangle contained in $\Gamma(R)$. $\square$

Lemma 4.9 *Let $(R, \mathfrak{m})$ be a finite local ring and $\Gamma(R) = K_3$. Then $R \cong \frac{\mathbb{Z}_2[x, y]}{(x^2, xy, y^2)}$ or $\frac{\mathbb{Z}_4[x]}{(2x, x^2)}$ or $R \cong \frac{\mathbb{F}_4[x]}{(x^2)}$ or $R \cong \frac{\mathbb{Z}_4[x]}{(x^2 + x + 1)}$. In particular, $\mathfrak{m}^2 = 0$.*

Proof. Among all rings in Theorem 4.2, there are exactly four rings $\frac{\mathbb{Z}_2[x, y]}{(x^2, xy, y^2)}$, $\frac{\mathbb{Z}_4[x]}{(2x, x^2)}$, $\frac{\mathbb{F}_4[x]}{(x^2)}$ and $\frac{\mathbb{Z}_4[x]}{(x^2 + x + 1)}$ satisfy the assumption. Moreover, it is easy to see that $\mathfrak{m}^2 = 0$ if R is one of the above three rings. $\square$

Lemma 4.10 *Let $R \cong R_1 \times \cdots \times R_k$, where R_i is a finite local ring for every i and $k \geq 3$. If $\gamma(\Gamma(R)^{(2)}) \leq 1$, then $k = 3$ and $R_i \cong \mathbb{Z}_2$ for every i .*

Proof. The first statement follows if we can show that $k = 4$ implies that $\gamma(\Gamma(R)^{(2)}) \geq 2$. For this, suppose that $k = 4$. Let $u_1 = (1, 0, 0, 0)$, $u_2 = (0, 1, 0, 0)$, $u_3 = (1, 1, 0, 0)$, $v_1 = (0, 0, 1, 0)$, $v_2 = (0, 0, 0, 1)$ and $v_3 = (0, 0, 1, 1)$; then $u_i v_j = 0$ for all i, j , so that $K_{3,3}$ is a subgraph of $\Gamma(R)$ it follows that $K_{6,6}$ is a subgraph of $\Gamma(R)^{(2)}$ by Example 2.6. Therefore, $\gamma(\Gamma(R)^{(2)}) \geq 2$ by Lemma 2.3.

To finish the proof, let $R \cong R_1 \times R_2 \times R_3$. Assume further that $|R_1| \leq |R_2| \leq |R_3|$ and $|R_3| \geq 3$. Let $u_1 = (1, 0, 0)$, $u_2 = (0, 1, 0)$, $u_3 = (1, 1, 0)$, $v_1 = (0, 0, 1)$, and $v_2 = (0, 0, 2)$; then $u_i v_j = 0$ for all i, j , so that $K_{2,3}$ is a subgraph of $\Gamma(R)$ it follows that $K_{4,6}$ is a subgraph of $\Gamma(R)^{(2)}$ by Proposition 2.7(d). Therefore, $\gamma(\Gamma(R)^{(2)}) \geq 2$ by Lemma 2.3, a contradiction. $\square$

There are some concrete examples for which $\gamma(\Gamma_I(R)) = 1$.

Example 4.11 *Let R be a finite ring and I be a proper nonzero ideal of R . Then the following hold:*

- (a) *Suppose that $R/I \cong \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$ and $|I| = 2$. Then $\gamma(\Gamma_I(R)) = 1$.*
- (b) *Suppose that $R/I \cong \mathbb{Z}_{16}$ and $|I| = 2$. Then $\gamma(\Gamma_I(R)) = 1$.*

(c) Suppose that $\Gamma(R/I) = P_3$, R/I is local and $|I| = 3$. Then $\gamma(\Gamma_I(R)) = 1$.

(d) Suppose that $R/I \cong \mathbb{Z}_9$ or $R/I \cong \frac{\mathbb{Z}_2[x]}{(x^3)}$ and $|I| = 3$. Then $\gamma(\Gamma_I(R)) = 1$.

Proof. (a) By assumption, $\omega(\Gamma(R/I)) = 3$, therefore $\Gamma_I(R)$ is not planar by [7, Theorem 7.2]. By Remark 3.3(c), the graph $\Gamma_I(R)$ and the graph $\Gamma_{0 \times 0 \times 0 \times \mathbb{Z}_2}(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2)$ are the same. Thus, we may assume that $R = \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$ and $I = 0 \times 0 \times 0 \times \mathbb{Z}_2$. Let $u_1 = (1, 0, 0, 0)$, $u_2 = (0, 1, 0, 0)$, $u_3 = (0, 0, 1, 0)$, $v_1 = (1, 0, 0, 1)$, $v_2 = (0, 1, 0, 1)$, $v_3 = (0, 0, 1, 1)$, $w_{11} = (0, 1, 1, 0)$, $w_{12} = (0, 1, 1, 1)$, $w_{21} = (1, 0, 1, 0)$, $w_{22} = (1, 0, 1, 1)$, $w_{31} = (1, 1, 0, 0)$ and $w_{32} = (1, 1, 0, 1)$. Let $G = \Gamma_I(R)$; then

$$V(G) = \{u_1, u_2, u_3, v_1, v_2, v_3, w_{11}, w_{12}, w_{21}, w_{22}, w_{31}, w_{32}\}$$

and

$$E(G) = \begin{aligned} & \{u_i u_j, v_i v_j, u_i v_j, v_i u_j \mid 1 \leq i < j \leq 3\} \\ & \cup \{u_i w_{ij}, v_i w_{ij} \mid i = 1, 2, 3, j = 1, 2\} \end{aligned}.$$

To draw G on a torus, we first draw a subgraph of G on a torus. Let G_1 be the subgraph of G induced by $\{u_1, u_2, u_3, v_1, v_2, v_3\}$; then Figure 3-1 gives the presentation of G_1 on a torus. To obtain a presentation of G , one can simply insert $\{w_{11}, w_{12}, u_1 w_{11}, u_1 w_{12}, v_1 w_{11}, v_1 w_{12}\}$ into the face F_1 , $\{w_{2j}, u_2 w_{2j}, v_2 w_{2j} \mid j = 1, 2\}$ into the face F_2 and $\{w_{3j}, u_3 w_{3j}, v_3 w_{3j} \mid j = 1, 2\}$ into the face F_3 .

(b) By assumption, $\Gamma(R/I)$ contains a triangle, therefore $\Gamma_I(R)$ is not planar by [7, Theorem 7.2]. By Remark 3.3(c), the graph $\Gamma_I(R)$ and the graph $\Gamma_{0 \times \mathbb{Z}_2}(\mathbb{Z}_{16} \times \mathbb{Z}_2)$ are the same. Thus, we may assume that $R = \mathbb{Z}_{16} \times \mathbb{Z}_2$ and $I = 0 \times \mathbb{Z}_2$. Let $u_1 = (4, 0)$, $u_2 = (8, 0)$, $u_3 = (12, 0)$, $v_1 = (4, 1)$, $v_2 = (8, 1)$, $v_3 = (12, 1)$, $w_1 = (2, 0)$, $w_2 = (6, 0)$, $w_3 = (10, 0)$, $w_4 = (14, 0)$, $w_5 = (2, 1)$, $w_6 = (6, 1)$, $w_7 = (10, 1)$ and $w_8 = (14, 1)$. Let $G = \Gamma_I(R)$; then

$$V(G) = \{u_1, u_2, u_3, v_1, v_2, v_3, w_1, \dots, w_8\}$$

and

$$E(G) = \begin{aligned} & \{u_i u_j, v_i v_j, u_i v_j, v_i u_j \mid 1 \leq i < j \leq 3\} \\ & \cup \{u_i v_i \mid i = 1, 2, 3\} \\ & \cup \{u_2 w_j, v_2 w_j \mid j = 1, \dots, 8\} \end{aligned}.$$

To draw G on a torus, we first draw a subgraph of G on a torus. Let G_1 be the subgraph of G induced by $\{u_1, u_2, u_3, v_1, v_2, v_3\}$; then Figure 3-2 gives the presentation of G_1 on a torus. To obtain a presentation of G , one can simply insert $\{w_j, u_2w_j, v_2w_j \mid j = 1, \dots, 8\}$ into the face F of Figure 3-2.




Figure 3-1




Figure 3-2

(c) By Theorem 4.2, $R/I \cong \mathbb{Z}_8$ or $R/I \cong \frac{\mathbb{Z}_2[x]}{(x^3)}$. Since the graph $\Gamma_I(R)$ can be viewed as the graph $\Gamma_{0 \times \mathbb{Z}_3}(R/I \times \mathbb{Z}_3)$ and the graphs $\Gamma_{0 \times \mathbb{Z}_3}(\mathbb{Z}_8 \times \mathbb{Z}_3)$ and $\Gamma_{0 \times \mathbb{Z}_3}(\frac{\mathbb{Z}_2[x]}{(x^3)} \times \mathbb{Z}_3)$ are the same, we may assume that $R = \mathbb{Z}_8 \times \mathbb{Z}_3$ and $I = 0 \times \mathbb{Z}_3$. Let $u_1 = (4, 0)$, $u_2 = (4, 1)$, $u_3 = (4, 2)$, $v_1 = (2, 0)$, $v_2 = (2, 1)$, $v_3 = (2, 2)$, $v_4 = (6, 0)$, $v_5 = (6, 1)$ and $v_6 = (6, 2)$. Let $G = \Gamma_I(R)$; then

$$V(G) = \{u_1, u_2, u_3, v_1, \dots, v_6\}$$

and

$$E(G) = \{u_i v_j \mid 1 \leq i \leq 3, 1 \leq j \leq 6\} \cup \{u_i u_j \mid 1 \leq i < j \leq 3\}.$$

Figure 4 gives the presentation of G on a torus.




Figure 4

(d) If $R/I \cong \mathbb{Z}_9$ or $R/I \cong \frac{\mathbb{Z}_2[x]}{(x^3)}$, then $a^2 = 0$ for every element in $\Gamma(R/I)$, so that $\Gamma_I(R) = K_6$, it follows that $\gamma(\Gamma_I(R)) = 1$. $\square$

To illustrate examples of $\gamma(\Gamma_I(R)) \geq 2$, we need a Lemma.

Lemma 4.12 *Let $(R, \mathfrak{m})$ be a finite local ring and I be a proper nonzero ideal of R . Suppose that the presentation of $\Gamma(R/I)$ is either G or G' and $\mathfrak{m}/I = \{0, u, v_1, \dots, v_6\}$ such that $u^2 = v_5^2 = v_6^2 = 0$. Then $\gamma(\Gamma_I(R)) \geq 2$.*

G




Figure 5-1

G'




Figure 5-2

Proof. We may assume that $|I| = 2$ and therefore the graph $\Gamma_I(R)$ can be viewed as the graph $\Gamma_{0 \times \mathbb{Z}_2}(R/I \times \mathbb{Z}_2)$. By abuse of notation, let $u = (u, 0)$ and $v_i = (v_i, 0)$ for $i = 1, \dots, 6$. Furthermore, let $u' = (u, 1)$ and $v'_i = (v_i, 1)$ for $i = 1, \dots, 6$. Then

$$V(\Gamma_I(R)) = \{u, u', v_1, \dots, v_6, v'_1, \dots, v'_6\}$$

and

$$\begin{aligned} E(\Gamma_I(R)) \supseteq & \{uu', uv_i, uv'_i, u'v_i, u'v'_i \mid i = 1, \dots, 6\} \\ & \cup \{v_5v'_5, v_5v_6, v_5v'_6, v'_5v_6, v'_5v'_6, v_6v'_6\} \\ & \cup \{v_i v_{i+1}, v_i v'_{i+1}, v'_i v_{i+1}, v'_i v'_{i+1} \mid i = 1, 3\} \end{aligned}$$

Let H be the subgraph of $\Gamma_I(R)$ such that $V(H) = V(\Gamma_I(R))$ and

$$\begin{aligned} E(H) = & \{uu', uv_i, uv'_i, u'v_i, u'v'_i \mid i = 1, \dots, 6\} \\ & \cup \{v_5v'_5, v_5v_6, v_5v'_6, v'_5v_6, v'_5v'_6, v_6v'_6\} \\ & \cup \{v_i v_{i+1}, v_i v'_{i+1}, v'_i v_{i+1}, v'_i v'_{i+1} \mid i = 1, 3\} \end{aligned}$$

Let $H_1 = H - \{v_5, v'_5, v_6, v'_6\}$ and H_2 be the subgraph induced by $\{v_5, v'_5, v_6, v'_6\}$; then there are edges $\{uv_5, uv'_5, uv_6, uv'_6\}$ of H join H_2 to H_1 and $H_2 = K_4$, it follows that $\gamma(H) > \gamma(H_1)$ by Lemma 2.8. Therefore, to finish the proof

it is enough to show that H_1 is not planar. For this, observe that H_1 has 10 vertices and 25 edges. If H_1 can be drawn in a plane, then H_1 has 17 faces by Euler characteristic formula. However, the edges and faced satisfies $50 \geq 17 \cdot 3$, a contradiction. Thus, $\gamma(H_1) \geq 1$. $\square$

Example 4.13 Let R be a finite ring and I be a proper nonzero ideal of R .

- (i) If R/I is one of the following local rings: (a) $\mathbb{Z}_4[x, y]/(x^2, y^2, xy - 2)$, (b) $\mathbb{Z}_2[x, y]/(x^2, y^2)$, (c) $\mathbb{Z}_4[x]/(x^2)$, then $\gamma(\Gamma_I(R)) \geq 2$.
(ii) If R/I is one of the following local rings: (a) $\mathbb{Z}_2[x, y]/(x^3, xy, y^2 - x^2)$, (b) $\mathbb{Z}_4[x]/(x^3, x^2 - 2x)$, (c) $\mathbb{Z}_4[x, y]/(x^3, x^2 - 2, xy, y^2 - 2)$, (d) $\mathbb{Z}_8[x]/(x^2 - 4, 2x)$, then $\gamma(\Gamma_I(R)) \geq 2$.

Proof. We may assume that $|I| = 2$.

(i) If R/I is one of the rings in (ii), then the graph of $\Gamma(R/I)$ is G' in Figure 5-1 by [5, Example 2.7]. Hence by Lemma 4.12, we need to find 3 elements in each ring with nilpotency 2. If $R = \mathbb{Z}_4[x, y]/(x^2, y^2, xy - 2)$, then $u = \bar{2}$, $v_5 = \bar{x} + \bar{y}$ and $v_6 = \bar{x} + \bar{y} + \bar{2}$ are the required 3 elements. If $R = \mathbb{Z}_2[x, y]/(x^2, y^2)$, then $u = \bar{x}\bar{y}$, $v_5 = \bar{x} + \bar{y}$ and $v_6 = \bar{x} + \bar{y} + \bar{x}\bar{y}$ are the required 3 elements. If $R = \mathbb{Z}_4[x]/(x^2)$, then $u = \bar{2}x$, $v_5 = \bar{2} + \bar{x}$ and $v_6 = \bar{2} + \bar{3}x$ are the required 3 elements.

(ii) If R/I is one of the ring in (i), then the graph of $\Gamma(R/I)$ is G' in Figure 5-2 by [5, Example 2.6]. Hence by Lemma 4.12, we need to find 3 elements in each ring with nilpotency 2. If $R = \mathbb{Z}_2[x, y]/(x^3, xy, y^2 - x^2)$, then $u = \bar{x}^2$, $v_5 = \bar{x} + \bar{y}$, and $v_6 = \bar{x} + \bar{y} + \bar{x}^2$ are the required 3 elements. If $R = \mathbb{Z}_4[x]/(x^3, x^2 - 2x)$, then $u = \bar{2}x$, $v_5 = \bar{2}$ and $v_6 = \bar{2} + \bar{2}x$ are the required 3 elements. If $R = \mathbb{Z}_4[x, y]/(x^3, x^2 - 2, xy, y^2 - 2)$, then $u = \bar{2}$, $v_5 = \bar{x} + \bar{y}$ and $v_6 = \bar{x} + \bar{y} + \bar{2}$ are the required 3 elements. If $R = \mathbb{Z}_8[x]/(x^2 - 4, 2x)$, then $u = \bar{4}$, $v_5 = \bar{2} + \bar{x}$ and $v_6 = \bar{6} + \bar{x}$ are the required 3 elements. $\square$

There are some sufficient conditions for which $\gamma(\Gamma_I(R)) \geq 2$.

Lemma 4.14 Let R be a finite ring and I be a proper nonzero ideal of R . Suppose that $\gamma(\Gamma(R/I)) \geq 2$. Then $\gamma(\Gamma_I(R)) \geq 2$.

Proof. The conclusion follows as

$$2 \leq \gamma(\Gamma(R/I)) \leq \gamma(\Gamma(R/I)^{(2)}) \leq \gamma(\Gamma_I(R)).$$

$\square$

Lemma 4.15 Let R be a finite ring and I be a proper nonzero ideal of R . Suppose the following hold:

(a) R/I is local with unique maximal ideal $\mathfrak{m}/I$.

(b) $|R/\mathfrak{m}| = 2$.

(c) $\gamma(\Gamma(R/I)) = 1$.

Then $\gamma(\Gamma_I(R)) \geq 2$.

Proof. By (a) and (c), R/I is isomorphic to one of the rings in Theorem 4.3. However, if we take (b) into account, then $K_{3,4}$ is a subgraph of every ring in Theorem 4.3 with (b) holds. Thus, $2 \leq \gamma(K_{6,8}) \leq \gamma(\Gamma(R/I)^{(2)}) \leq \gamma(\Gamma_I(R))$. $\square$

Theorem 4.16 *Let R be a finite ring and I be a proper nonzero ideal of R . Suppose that $\omega(\Gamma(R/I)) \geq 4$ or $\gamma(\Gamma(R/I)) \geq 1$. Then $\gamma(\Gamma_I(R)) \geq 2$.*

Proof. Since R is finite, $R/I \cong R_1 \times \cdots \times R_k$, where R_i is a finite local ring for every i . If $k \geq 4$, then by Lemma 4.10 and the facts that $|I| \geq 2$ and $\Gamma((R/I)^{(2)})$ is a subgraph of $\Gamma_I(R)$, we see that $\gamma(\Gamma_I(R)) \geq 2$. Therefore, we may assume that $k \leq 2$ or $R/I \cong \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$. However, $R \cong \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$ implies that $\omega(\Gamma(R/I)) = 3$ and $\Gamma(R/I)$ is planar, a contradiction. Thus, $k \leq 2$.

Now, suppose that R/I is not local and $R/I \cong R_1 \times R_2$ with $|R_1| \leq |R_2|$. If $|R_1| \geq 3$ and $|R_2| \geq 4$, then $K_{2,3}$ is a subgraph of $\Gamma(R/I)$, so that $\gamma(\Gamma(R/I)^{(2)}) \geq 2$ by Proposition 2.7(d), it follows that $\gamma(\Gamma_I(R)) \geq 2$. If $|R_1| = |R_2| = 3$, then $R/I \cong \mathbb{Z}_3 \times \mathbb{Z}_3$, which contradicts to the assumptions that $\omega(\Gamma(R/I)) \geq 4$ or $\gamma(\Gamma(R/I)) \geq 1$. So, we may assume that $R_1 \cong \mathbb{Z}_2$. Since $\gamma(\Gamma(\mathbb{Z}_2 \times R_2)) \geq 1$ or $\omega(\Gamma(\mathbb{Z}_2 \times R_2)) \geq 4$, by Lemma 4.8, R_2 is neither a field nor satisfies $|\Gamma(R_2)| = 1$. Therefore $|\Gamma(R_2)| \geq 2$. However, by Lemma 4.8 again $K_{2,3}$ is a subgraph of $\Gamma(R/I)$, it follows that $\gamma(\Gamma(R/I)^{(2)}) \geq 2$ by Proposition 2.7(d). Hence, $\gamma(\Gamma_I(R)) \geq 2$.

Finally we assume that R/I is local with unique maximal ideal $\mathfrak{m}/I$.

Case 1. $(\mathfrak{m}/I)^2 = 0$. In this case, $\omega(\Gamma(R/I)) \geq 4$ or $\gamma(\Gamma(R/I)) \geq 1$ implies that there are 4 distinct nonzero zero-divisors $u_1, \dots, u_4$ in R/I such that $u_i u_j = 0$ for all i, j , so that K_8 is a subgraph of $\Gamma(R/I)^{(2)}$, it follows that $\gamma(\Gamma_I(R)) \geq 2$.

Case 2. $(\mathfrak{m}/I)^2 \neq 0$. In this case, observe $\Gamma(R/I)^{(2)}$ is a subgraph of $\Gamma_I(R)$. Hence, by Lemma 4.6, we may assume that $|R/\mathfrak{m}| = 2$ as $|R/\mathfrak{m}| \geq 3$ implies that $\gamma(\Gamma_I(R)) \geq 2$. Therefore, $|\Gamma(R/I)| = 2^n - 1$ for some positive

integer n . Observe that $|\Gamma(R/I)|$ is 1 or 3 are not possible. Moreover, the assumption that $\gamma(\Gamma(R/I)) \geq 1$ implies that $\gamma(\Gamma_I(R)) \geq 2$ by Lemma 4.14 and Lemma 4.15. Therefore, we may assume finally that $\omega(\Gamma(R/I)) \geq 4$ and $\Gamma(R/I)$ is planar. However, if so, then by Theorem 4.2, $|\Gamma(R/I)| = 7$ and R/I is isomorphic to $\mathbb{Z}_{16}$ or isomorphic to one of the rings in Example 4.13, so that $\omega(\Gamma(S)) = 3$, a contradiction. The proof is now complete. $\square$

Theorem 4.17 *Let R be a finite ring and I be a proper nonzero ideal of R which is not a prime ideal. Suppose that $\omega(\Gamma(R/I)) \leq 2$. Then $\gamma(\Gamma_I(R)) \leq 1$ if and only if one of the following holds:*

- (a) $R/I \cong \mathbb{Z}_3 \times \mathbb{Z}_3$ and $|I| \leq 2$.
- (b) $R/I \cong \mathbb{Z}_2 \times \mathbb{Z}_2$ and $|I| \leq 4$.
- (c) $R/I \cong \mathbb{Z}_2 \times \mathbb{Z}_3$ and $|I| \leq 3$.
- (d) $R/I \cong \mathbb{Z}_2 \times \mathbb{Z}_4$ and $|I| \leq 2$.
- (e) $R/I \cong \mathbb{Z}_2 \times \frac{\mathbb{Z}_2[x]}{(x^2)}$ and $|I| \leq 2$.
- (f) $R/I \cong \mathbb{Z}_2 \times \mathbb{F}_q$ with $q \geq 4$ and $|I| \leq 2$.
- (g) $R/I = \mathbb{Z}_4$ and $|I| \leq 7$.
- (h) $R/I = \frac{\mathbb{Z}_2[x]}{(x^2)}$ and $|I| \leq 7$.
- (i) $R/I = \mathbb{Z}_9$ and $|I| \leq 3$.
- (j) $R/I = \frac{\mathbb{Z}_3[x]}{(x^2)}$ and $|I| \leq 3$.
- (k) $R/I = \mathbb{Z}_8$ and $|I| \leq 3$.
- (l) $R/I = \frac{\mathbb{Z}_2[x]}{(x^3)}$ and $|I| \leq 3$.
- (m) $R/I = \frac{\mathbb{Z}_4[x]}{(x^2-2, x^3)}$ and $|I| \leq 3$.

Proof. ($\Rightarrow$) Assume that $\gamma(\Gamma_I(R)) \leq 1$. From the first paragraph of the proof of Theorem 4.16, we may assume that $R/I = \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$ or R/I is local or R/I is a product of two finite local rings. However, $R/I = \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$

implies that $\omega(\Gamma(R/I)) = 3$, a contradiction. Thus, R/I is local or a product of two finite local rings.

Now, suppose that R/I is not local and $R/I \cong R_1 \times R_2$ with $|R_1| \leq |R_2|$. From the second paragraph of the proof of Theorem 4.16, we may assume that $R/I \cong \mathbb{Z}_3 \times \mathbb{Z}_3$ or $R_1 \cong \mathbb{Z}_2$. If $R/I \cong \mathbb{Z}_3 \times \mathbb{Z}_3$, then $\Gamma(R/I) = C_4 = K_{2,2}$ and R/I has no nilpotent, so that $|I| \leq 2$. Now, we assume that $R_1 = \mathbb{Z}_2$. By Lemma 4.8, R_2 is either a field or satisfies $|\Gamma(R_2)| = 1$. It follows that R/I is isomorphic to one of the following rings: $\mathbb{Z}_2 \times \mathbb{Z}_2$, $\mathbb{Z}_2 \times \mathbb{Z}_3$, $\mathbb{Z}_2 \times \mathbb{Z}_4$, $\mathbb{Z}_2 \times \frac{\mathbb{Z}_2[x]}{(x^2)}$ and $\mathbb{Z}_2 \times \mathbb{F}_q$, where $q \geq 4$.

If $R/I \cong \mathbb{Z}_2 \times \mathbb{Z}_2$, then $\Gamma(R/I) = K_2$. Since R/I has no nilpotent, $\Gamma_I(R) = K_{t,t}$ if $|I| = t$, it follows that $|I| \leq 4$ by Lemma 2.3.

If $R/I \cong \mathbb{Z}_2 \times \mathbb{Z}_3$, then $\Gamma(R/I) = K_{1,2}$. Since R/I has no nilpotent, $\Gamma_I(R) = K_{t,2t}$ if $|I| = t$, it follows that $|I| \leq 3$ by Lemma 2.3.

If R/I is isomorphic to $\mathbb{Z}_2 \times \mathbb{Z}_4$ or $\mathbb{Z}_2 \times \frac{\mathbb{Z}_2[x]}{(x^2)}$ or $\mathbb{Z}_2 \times \mathbb{F}_q$ with $q \geq 4$, then $K_{1,3}$ is a subgraph of $\Gamma(R/I)$, it follows that $|I| \leq 2$ by Lemma 2.3.

From the above, we can finally assume that R/I is local with unique maximal ideal $\mathfrak{m}/I$. By Theorem 4.16, $\Gamma(R/I)$ is planar. So we only consider the rings in Theorem 4.2.

Case 1. $(\mathfrak{m}/I)^2 = 0$. In this case, $\Gamma(R/I) \cong K_t$ if $|(\mathfrak{m}/I)| = t + 1$, so that $\Gamma(R/I)$ is K_1 or K_2 as $\omega(\Gamma(R/I)) \leq 2$, it follows that R/I is isomorphic to the following rings: $\mathbb{Z}_4$, $\frac{\mathbb{Z}_2[x]}{(x^2)}$, $\mathbb{Z}_9$ and $\frac{\mathbb{Z}_3[x]}{(x^2)}$.

If R/I is isomorphic to $\mathbb{Z}_4$ or $\frac{\mathbb{Z}_2[x]}{(x^2)}$, then $\Gamma_I(R) = K_t$ if $|I| = t$, it follows that $|I| \leq 7$ by Lemma 2.2.

If R/I is isomorphic to $\mathbb{Z}_9$ or $\frac{\mathbb{Z}_3[x]}{(x^2)}$, then $\Gamma_I(R) = K_{2t}$ if $|I| = t$, it follows that $|I| \leq 3$ by Lemma 2.2.

Case 2. $(\mathfrak{m}/I)^2 \neq 0$. In this case, $|R/\mathfrak{m}| = 2$ as $|R/\mathfrak{m}| \geq 3$ implies that $\gamma(\Gamma_I(R)) \geq 2$ ($\Gamma(R/I)^{(2)}$ is a subgraph of $\Gamma_I(R)$) by Lemma 4.6. Moreover, by Theorem 4.16, $\Gamma(R/I)$ is planar. Hence by Theorem 4.2, $|\Gamma(R/I)| = 1, 3, 7$. From the above, we see that $|\Gamma(R/I)| = 3, 7$. However, if $|\Gamma(R/I)| = 7$, then R/I is isomorphic to $\mathbb{Z}_{16}$ or isomorphic to one of the rings in Example 4.13, so that $\Gamma(R/I)$ contains a triangle, a contradiction. Thus, $|\Gamma(R/I)| = 3$ and $|\Gamma(R/I)|$ is P_3 as $\Gamma(R/I)$ contains no triangles and is connected. Therefore, R/I is isomorphic to the following rings: $\mathbb{Z}_8$, $\frac{\mathbb{Z}_2[x]}{(x^3)}$ and $\frac{\mathbb{Z}_4[x]}{(x^2-2, x^3)}$. In either cases, $K_{t,2t}$ is a subgraph of $\Gamma_I(R)$ if $|I| = t$, therefore $t \leq 3$.

($\Leftarrow$) Assume that $R/I \cong \mathbb{Z}_3 \times \mathbb{Z}_3$ and $|I| \leq 2$. Then $\Gamma(R/I) = K_{2,2}$ and R/I has no nilpotent. It follows that $\Gamma_I(R) = K_{2t,2t}$ with $|I| = t \leq 2$.

Therefore, $\gamma(\Gamma(R/I)) \leq 1$ by Lemma 2.3.

Assume that $R/I \cong \mathbb{Z}_2 \times \mathbb{Z}_2$ and $|I| \leq 4$. Then $\Gamma(R/I) = K_2$ and R/I has no nilpotent. It follows that $\Gamma_I(R) = K_{2t}$ with $|I| = t \leq 4$. Therefore, $\gamma(\Gamma(R/I)) \leq 1$ by Lemma 2.3.

Assume that $R/I \cong \mathbb{Z}_2 \times \mathbb{Z}_3$ and $|I| \leq 3$. Then $\Gamma(R/I) = K_{1,2}$ and R/I has no nilpotent. It follows that $\Gamma_I(R) = K_{t,2t}$ with $|I| = t \leq 3$. Therefore, $\gamma(\Gamma(R/I)) \leq 1$ by Lemma 2.3.

Assume that $R \cong \mathbb{Z}_2 \times \mathbb{Z}_4$ or $R \cong \mathbb{Z}_2 \times \frac{\mathbb{Z}_2[x]}{(x^2)}$ or $R/I \cong \mathbb{Z}_2 \times \mathbb{F}_q$ with $q \geq 4$ and $|I| \leq 2$. Then $\Gamma_I(R)$ is planar by [7, Theorem 7.2].

Assume that $R \cong \mathbb{Z}_4$ or $R \cong \frac{\mathbb{Z}_2[x]}{(x^2)}$. In either cases, $\Gamma_I(R) = K_t$ if $|I| = t$. Therefore if $|I| \leq 7$, then $\gamma(\Gamma_I(R)) \leq 1$ by Lemma 2.2.

Assume that $R/I \cong \mathbb{Z}_9$ or $R/I \cong \mathbb{Z}_3[x]/(x^2)$. Then for every zero-divisor $a \in R/I$ satisfies $a^2 = 0$. Therefore, $\Gamma_I(R) = K_{2t}$ if $|I| = t$. Thus, if $|I| \leq 3$, then $\gamma(\Gamma_I(R)) \leq 1$ by Lemma 2.2.

Assume that R/I is isomorphic to the following rings: $\mathbb{Z}_8$, $\frac{\mathbb{Z}_2[x]}{(x^3)}$ and $\frac{\mathbb{Z}_4[x]}{(x^2-2, x^3)}$. In either cases, if $|I| \leq 3$ then $\gamma(\Gamma_I(R)) \leq 1$ by Example 4.11(c). $\square$

Theorem 4.18 *Let R be a finite ring and I be a proper nonzero ideal of R . Suppose that $\omega(\Gamma(R/I)) = 3$. Then $\gamma(\Gamma_I(R)) = 1$ if and only if $|I| = 2$ and R/I is isomorphic to one of the following:*

(a) $\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$.

(b) $\mathbb{Z}_{16}$.

(c) $\frac{\mathbb{Z}_2[x, y]}{(x^2, xy, y^2)}$.

(d) $\frac{\mathbb{Z}_4[x]}{(2x, x^2)}$.

(e) $\frac{\mathbb{F}_4[x]}{(x^2)}$.

(f) $\frac{\mathbb{Z}_4[x]}{(x^2 + x + 1)}$.

Proof. ($\Rightarrow$) Assume that $\gamma(\Gamma_I(R)) = 1$. From the first paragraph of the proof of Theorem 4.16, we may assume that $R/I \cong \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$ or R/I is local

or R/I is a product of two finite local rings.

Assume that $R/I \cong \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$. If $|I| \geq 3$, then $\gamma(\Gamma(R/I)^{(3)}) \geq 2$ as $K_{1,3}$ is a subgraph of $\Gamma(R/I)$, it follows that $\gamma(\Gamma_I(R)) \geq 2$. Moreover, If $|I| = 2$, then $\gamma(\Gamma_I(R)) = 1$ by Example 4.11(a).

Assume that R/I is not local and $R/I \cong R_1 \times R_2$ with $|R_1| \leq |R_2|$. From the second paragraph of the proof of Theorem 4.16, we may assume that $R/I \cong \mathbb{Z}_3 \times \mathbb{Z}_3$ or $R_1 \cong \mathbb{Z}_2$. However, if $R/I \cong \mathbb{Z}_3 \times \mathbb{Z}_3$, then $\omega(\Gamma(R/I)) = 2$, a contradiction. Thus, $R_1 \cong \mathbb{Z}_2$. Moreover, if R_2 is a field or $\Gamma(R_2)$ is a point then $\Gamma(R/I)$ contains no cycles by Lemma 4.8 which contradicts to the assumption that $\omega(\Gamma(R/I)) = 3$. If $|\Gamma(R_2)| \geq 2$, then $K_{2,3}$ is a subgraph of $\Gamma(R/I)$, it follows that $\gamma(\Gamma_I(R)) \geq 2$, a contradiction again.

From the above, we can finally assume that R/I is local with unique maximal ideal $\mathfrak{m}/I$. Observe that the assumption $\omega(\Gamma(R/I)) = 3$ implies that $|(\mathfrak{m}/I)| \geq 4$.

Case 1. $(\mathfrak{m}/I)^2 = 0$. In this case, $\Gamma(R/I) \cong K_t$ if $|(\mathfrak{m}/I)| = t + 1$, so that $\Gamma(R/I) \cong K_3$ as $\omega(\Gamma(R/I)) = 3$, it follows that R/I is isomorphic to $\frac{\mathbb{Z}_2[x, y]}{(x^2, xy, y^2)}$ or $\frac{\mathbb{Z}_4[x]}{(2x, x^2)}$ or $\frac{\mathbb{F}_4[x]}{(x^2)}$ or $\frac{\mathbb{Z}_4[x]}{(x^2 + x + 1)}$ by Lemma 4.9. Moreover, every element $a \in R/I$ satisfies $a^2 = 0$. Thus, $\Gamma_I(R) = K_{3t}$ if $|I| = t$. It follows that $\gamma(\Gamma_I(R)) = 1$ if and only if $|I| = 2$.

Case 2. $(\mathfrak{m}/I)^2 \neq 0$. In this case, $|R/\mathfrak{m}| = 2$ as $|R/\mathfrak{m}| \geq 3$ implies that $\gamma(\Gamma_I(R)) \geq 2$ ($\Gamma(R/I)^{(2)}$ is a subgraph of $\Gamma_I(R)$) by Lemma 4.6. Moreover, by Theorem 4.16, $\Gamma(R/I)$ is planar. Hence by Theorem 4.2, $|\Gamma(R/I)| = 1, 3, 7$. From the above, we may assume that $|\Gamma(R/I)| = 7$. However, if so R/I is isomorphic to $\mathbb{Z}_{16}$ or isomorphic to one of the rings in Example 4.13. Since $\gamma(\Gamma_I(R)) \geq 2$ if R/I is isomorphic to one of the rings in Example 4.13, we see that $R/I \cong \mathbb{Z}_{16}$. If $|I| \geq 3$, then $\gamma(\Gamma_I(R)) \geq 2$ as $K_{1,3}$ is a subgraph of $\Gamma(R/I)$. Thus $|I| = 2$.

($\Leftarrow$) Assume that R/I is isomorphic to one of the following 2 rings: $\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$ and $\mathbb{Z}_{16}$. If $|I| = 2$, then $\gamma(\Gamma_I(R)) = 1$ by Example 4.11.

Assume that R/I is isomorphic to one of the following 4 rings: $\frac{\mathbb{Z}_2[x, y]}{(x^2, xy, y^2)}$, $\frac{\mathbb{Z}_4[x]}{(2x, x^2)}$, $\frac{\mathbb{F}_4[x]}{(x^2)}$ and $\frac{\mathbb{Z}_4[x]}{(x^2 + x + 1)}$. If $|I| = 2$, then $\gamma(\Gamma_I(R)) = 1$ as $\Gamma_I(R) = K_6$. $\square$

References

- [1] S. Akbari, H.R. Maimani and S. Yassemi, *When a zero-divisor graph is planar or a complete r -partite graph*, J. Algebra **270** (2003), 169–180.
- [2] S. Akbari and A. Mohammadian, *On the zero-divisor graph of a commutative ring*, J. Algebra **274** (2004), 847–855.
- [3] D.F. Anderson, A. Frazier, A. Lauve and P.S. Livingston, *The zero-divisor graph of a commutative ring, II*, Lecture Notes in Pure and Appl. Math. **220** (2001), 61–72.
- [4] D.F. Anderson and P.S. Livingston, *The zero-divisor graph of a commutative ring*, J. Algebra **217** (1999), 434–447.
- [5] H.-J. Chiang-Hsieh and H.-J. Wang, *Commutative rings with toroidal zero-divisor graphs*”, preprint.
- [6] H.R. Maimani, M.R. Pournaki and S. Yassemi, *Zero-divisor graph with respect to an ideal*, Comm. Algebra **34** (2006), no. 3, 923–929.
- [7] S. Redmond, *An ideal-based zero-divisor graph of a commutative ring*, Comm. Algebra **31** (2003), no. 9, 4425–4443.
- [8] G. Ringel and J.W.T. Youngs, *Solution of the Heawood map-coloring problem*, Proc. Nat. Acad. Sci. USA **60** (1968), 438–445.
- [9] N.O. Smith, *Planar zero-divisor graphs*, International J. Commutative Rings **2** (2003), no. 4, 177–188.
- [10] H.-J. Wang, *Zero-divisor graphs of genus one*, J. Algebra **304** (2006), no. 2, 666–678.